List of Trend Tables		14. Low-birthweight live births, according to race and Hispanic origin of mother, geographic division, and State: United States, average annual 1992–94, 1995–97, and	
Health Status and Determinants		1998–2000	96
Population		Hispanic origin of mother, geographic division, and State: United States, average annual 1992–94, 1995–97, and 1998–2000	98
Resident population, according to age, sex, race, and Hispanic origin: United States, selected years 1950–2000	79	16. Legal abortion ratios , according to selected patient characteristics: United States, selected years 1973–98	100
Persons and families below poverty level, according to selected characteristics, race, and Hispanic origin:		17. Legal abortions , according to selected characteristics: United States, selected years 1973–98	101
United States, selected years 1973–2000	81	18. Contraceptive use among women 15–44 years of age, according to age, race, Hispanic origin, and method of	
Fertility and Natality		contraception: United States, 1982, 1988, and 1995	102
3. Crude birth rates, fertility rates , and birth rates by age of mother, according to race and Hispanic origin: United States, selected years 1950–2000	83	year of baby's birth, according to selected characteristics of mother: United States, average annual 1972–74 to 1993–94.	104
4. Women 15–44 years of age who have not had at least 1 live birth, by age: United States, selected years 1960–2001	85	Mortality	
5. Live births , according to detailed race and Hispanic origin of mother: United States, selected years		20. Infant, neonatal, and postneonatal mortality rates, according to detailed race and Hispanic origin of mother: United States, selected years 1983–99	105
1970–2000	86	21. Infant mortality rates for mothers 20 years of age and over, according to mother's education, detailed race, and Hispanic origin: United States, selected years 1983–99	107
7. Early prenatal care according to race and Hispanic origin of mother, geographic division, and State:	67	22. Infant mortality rates according to birthweight: United States, selected years 1983–99	108
United States, average annual 1992–94, 1995–97, and 1998–2000	88	23. Infant mortality rates , fetal mortality rates, and perinatal mortality rates, according to race: United States, selected years 1950–99	109
Hispanic origin of mother: United States, selected years 1970–2000	90	24. Infant mortality rates , according to race, Hispanic origin, geographic division, and State: United States, average annual 1989–91 and 1997–99	110
9. Nonmarital childbearing according to detailed race and Hispanic origin of mother, and maternal age and birth rates for unmarried women by race and Hispanic origin of mother: United States, selected years 1970–2000	91	25. Neonatal mortality rates , according to race, Hispanic origin, geographic division, and State: United States, average annual 1989–91 and 1997–99	112
10. Maternal education for live births, according to detailed race and Hispanic origin of mother: United States, selected	00	26. Infant mortality rates and international rankings: Selected countries, selected years 1960–98	114
years 1970–2000	92	27. Life expectancy at birth and at 65 years of age, according to sex: Selected countries, 1992 and 1997	115
and education: Selected States, 1989–2000	93	28. Life expectancy at birth, at 65 years of age, and at 75 years of age, according to race and sex: United States, selected years 1900–99	116
detailed race, Hispanic origin, and smoking status: United States, selected years 1970–2000	94	29. Age-adjusted death rates , according to race, Hispanic origin, geographic division, and State: United States,	
13. Low-birthweight live births among mothers 20 years of age and over, by mother's detailed race, Hispanic origin, and education: United States, 1989–2000	95	average annual 1979–81, 1989–91, and 1997–99	117

31. Years of potential life lost before age 75 for selected causes of death, according to sex, race, and Hispanic origin: United States, selected years 1980–99	123	48. Death rates for firearm-related injuries , according to sex, race, Hispanic origin, and age: United States, selected years 1970–99	172
32. Leading causes of death and numbers of deaths, according to sex, race, and Hispanic origin: United States, 1980 and 1999	127	49. Deaths from selected occupational diseases for persons 15 years of age and over: United States, selected years 1980–99	175
33. Leading causes of death and numbers of deaths, according to age: United States, 1980 and 1999	131	50. Occupational injury deaths and rates by industry, sex, age, race, and Hispanic origin: United States, 1992–2000	176
34. Age-adjusted death rates, according to race, sex, region, and urbanization level: United States, average annual 1984–86, 1989–91, and 1997–99	133	Determinants and Measures of Health	
35. Age-adjusted death rates for persons 25–64 years of age for selected causes of death, according to sex and educational attainment: Selected States, 1994–99	136	51. Occupational injuries with lost workdays in the private sector, according to industry: United States, selected years 1980–2000	178
36. Death rates for all causes, according to sex, race, Hispanic origin, and age: United States, selected years 1950–99	137	52. Healthy People 2010 Leading Health Indicators with 22 measures: United States, 1990–2000 and 2010 target .	179
37. Death rates for diseases of heart , according to sex, race, Hispanic origin, and age: United States, selected		53. Selected notifiable disease rates , according to disease: United States, selected years 1950–2000	181
years 1950–99	141	54. Acquired immunodeficiency syndrome (AIDS) cases, according to age at diagnosis, sex, detailed race, and Hispanic origin: United States, selected years 1985–2001	182
to sex, race, Hispanic origin, and age: United States, selected years 1950–99	144	55. Acquired immunodeficiency syndrome (AIDS) cases,	102
39. Death rates for malignant neoplasms , according to sex, race, Hispanic origin, and age: United States, selected years 1950–99	147	according to race, Hispanic origin, sex, and transmission category for persons 13 years of age and over at diagnosis: United States, selected years 1985–2001	183
40. Death rates for malignant neoplasms of trachea, bronchus, and lung, according to sex, race, Hispanic origin, and age: United States, selected years 1950–99	151	56. Age-adjusted cancer incidence rates for selected cancer sites, according to sex, race, and Hispanic origin: Selected geographic areas, 1990–98	185
41. Death rates for malignant neoplasm of breast for females, according to race, Hispanic origin, and age: United States, selected years 1950–99	154	57. Five-year relative cancer survival rates for selected cancer sites, according to race and sex: Selected geographic areas, 1974–79, 1980–82, 1983–85, 1986–88,	100
42. Death rates for chronic lower respiratory diseases , according to sex, race, Hispanic origin, and age: United States, selected years 1980–99	156	and 1989–97	188
43. Death rates for human immunodeficiency virus (HIV)		selected years 1997–2000	189
disease, according to sex, race, Hispanic origin, and age: United States, selected years 1987–99	159	selected characteristics: United States, selected years 1991–2000	192
44. Maternal mortality for complications of pregnancy, childbirth, and the puerperium, according to race, Hispanic origin, and age: United States, selected years 1950–99	161	60. Suicidal ideation, suicide attempts, and injurious suicide attempts among students in grades 9–12, by sex,	
45. Death rates for motor vehicle-related injuries , according to sex, race, Hispanic origin, and age:		grade level, race, and Hispanic origin: United States, selected years 1991–2001	194
United States, selected years 1950–99	162	61. Current cigarette smoking by persons 18 years of age and over according to sex, race, and age: United States,	400
Hispanic origin, and age: United States, selected years 1950–99	166	selected years 1965–2000	196
47. Death rates for suicide , according to sex, race, Hispanic origin, and age: United States, selected years		by persons 25 years of age and over, according to sex, race, and education: United States, selected years 1974–2000	198
1950–99	169	13/4-2000	190

63. Current cigarette smoking by adults according to sex, race, Hispanic origin, age, and education: United States, average annual 1990–92, 1993–95, and 1999–2000	199	76. No usual source of health care among children under 18 years of age, according to selected characteristics: United States, average annual 1993–94, 1997–98, and	000
64. Use of selected substances in the past month by persons 12 years of age and over, according to age, sex, race, and Hispanic origin: United States, 1999–2000 65. Use of selected substances by high school seniors	201	77. Emergency department visits within the past 12 months among children under 18 years of age, according to selected characteristics: United States, selected years	226
and eighth-graders, according to sex and race: United States, selected years 1980–2001	203	78. No usual source of health care among adults 18–64	228
66. Cocaine-related emergency department episodes, according to age, sex, race, and Hispanic origin: United States, selected years 1990–2000	205	years of age, according to selected characteristics: United States, average annual 1993–94, 1995–96, 1997–98, and 1999–2000	231
67. Alcohol consumption by persons 18 years of age and over, according to sex, age, race, and Hispanic origin: United States, selected years 1997–2000	207	79. Emergency department visits within the past 12 months among adults 18 years of age and over, according to selected characteristics: United States, selected years 1997–2000.	233
68. Hypertension among persons 20 years of age and over, according to sex, age, race, and Hispanic origin: United States, 1960–62, 1971–74, 1976–80, and		80. Dental visits in the past year according to selected characteristics: United States, selected years 1997–2000	235
1988–94	210	81. Untreated dental caries according to age, sex, race and Hispanic origin, and poverty status: United States, 1971–74, 1982–84, and 1988–94	237
age and over, according to sex, age, race, and Hispanic origin: United States, 1960–62, 1971–74, 1976–80, and 1988–94	211	82. Use of mammography for women 40 years of age and over according to selected characteristics: United States, selected years 1987–2000	238
70. Healthy weight, overweight, and obesity among persons 20 years of age and over, according to sex, age, race, and Hispanic origin: United States, 1960–62, 1971–74, 1976–80, and 1988–94	213	83. Ambulatory care visits to physician offices and hospital outpatient and emergency departments by selected characteristics: United States, selected years 1995–2000	240
71. Overweight children and adolescents 6–19 years of age, according to sex, age, race, and Hispanic origin: United States, selected years 1963–65 through		84. Injury-related visits to hospital emergency departments by sex, age, and intent and mechanism of injury: United States, average annual 1995–96, 1997–98, and 1999–2000	242
Utilization of Health Resources	215	85. Ambulatory care visits to primary care and specialist physicians, according to selected characteristics and type of physician: United States, 1980, 1990, and 2000	244
Ambulatory Care		86. Substance abuse clients in specialty treatment units according to substance abused, geographic division, and State: United States, 1997–2000	246
72. Health care visits to doctor's offices, emergency departments, and home visits within the past 12 months, according to selected characteristics: United States,		87. Additions to mental health organizations according to type of service and organization: United States, selected years 1986–98	247
selected years 1997–2000	217	88. Home health care patients , according to age, sex, and diagnosis: United States, selected years 1992–2000	248
selected diseases, according to race, Hispanic origin, poverty status, and residence in metropolitan statistical area (MSA): United States, 1995–2000	220	89. Hospice patients , according to age, sex, and diagnosis: United States, selected years 1992–2000	249
74. Vaccination coverage among children 19–35 months of age according to geographic division, State, and selected		Inpatient Care	
urban areas: United States, 1995–2000	222	90. Discharges , days of care, and average length of stay in short-stay hospitals, according to selected characteristics: United States, selected years 1997–2000	250
according to selected characteristics: United States, average annual 1997–98 and 1999–2000	224	91. Discharges , days of care, and average length of stay in non-Federal short-stay hospitals, according to selected characteristics: United States, selected years 1980–2000	253

92. Discharges , days of care, and average length of stay in non-Federal short-stay hospitals for discharges with the diagnosis of human immunodeficiency virus (HIV) and for all discharges: United States, selected years 1986–2000	255	106. First-year and total enrollment of women in schools for selected health occupations, according to detailed race and Hispanic origin: United States, academic years 1971–72, 1980–81, 1990–91, and 1999–2000	278
93. Rates of discharges and days of care in non-Federal short-stay hospitals, according to sex, age, and selected first-listed diagnoses: United States, selected years		Facilities	
1990–2000	256	107. Hospitals , beds, and occupancy rates, according to type of ownership and size of hospital: United States, selected years 1975–2000	279
short-stay hospitals, according to sex, age, and selected first-listed diagnoses: United States, selected years 1990–2000	259	108. Mental health organizations and beds for 24-hour hospital and residential treatment according to type of organization: United States, selected years 1986–98	280
95. Ambulatory and inpatient procedures according to place, sex, age, and type of procedure: United States, selected years 1994–2000	262	109. Community hospital beds and average annual percent change, according to geographic division and State: United States, selected years 1940–2000	281
96. Hospital admissions , average length of stay, and outpatient visits, according to type of ownership and size of hospital, and percent outpatient surgery: United States, selected years 1975–2000	265	110. Occupancy rates in community hospitals and average annual percent change, according to geographic division and State: United States, selected years 1940–2000	283
97. Nursing home residents 65 years of age and over, according to age, sex, and race: United States, 1973–74, 1985, 1995, and 1999	266	111. Nursing homes , beds, occupancy, and residents, according to geographic division and State: United States, 1995–2000	284
98. Nursing home residents 65 years of age and over, according to selected functional status and age, sex, and race: United States, 1985, 1995, and 1999	267	Health Care Expenditures and Health Insurance	
Health Care Resources		National Health Expenditures	
Personnel 99. Persons employed in health service sites:			287
Personnel	269	National Health Expenditures 112. Total health expenditures as a percent of gross domestic product and per capita health expenditures in dollars: Selected countries and years 1960–99	287
Personnel 99. Persons employed in health service sites: United States, selected years 1970–2001 100. Active non-Federal physicians and doctors of medicine in patient care, according to geographic division		National Health Expenditures 112. Total health expenditures as a percent of gross domestic product and per capita health expenditures in dollars: Selected countries and years 1960–99	
Personnel 99. Persons employed in health service sites: United States, selected years 1970–2001	270	National Health Expenditures 112. Total health expenditures as a percent of gross domestic product and per capita health expenditures in dollars: Selected countries and years 1960–99	288
Personnel 99. Persons employed in health service sites: United States, selected years 1970–2001 100. Active non-Federal physicians and doctors of medicine in patient care, according to geographic division and State: United States, 1975, 1985, 1995, and 2000 101. Physicians, according to activity and place of medical education: United States and outlying U.S. areas, selected years 1975–2000 102. Primary care doctors of medicine, according to specialty: United States and outlying U.S. areas, selected years 1949–2000 103. Active health personnel according to occupation: United States, selected years 1980–2000	270 272	National Health Expenditures 112. Total health expenditures as a percent of gross domestic product and per capita health expenditures in dollars: Selected countries and years 1960–99	288 289 290
Personnel 99. Persons employed in health service sites: United States, selected years 1970–2001 100. Active non-Federal physicians and doctors of medicine in patient care, according to geographic division and State: United States, 1975, 1985, 1995, and 2000 101. Physicians, according to activity and place of medical education: United States and outlying U.S. areas, selected years 1975–2000 102. Primary care doctors of medicine, according to specialty: United States and outlying U.S. areas, selected years 1949–2000 103. Active health personnel according to occupation:	270 272 273	National Health Expenditures 112. Total health expenditures as a percent of gross domestic product and per capita health expenditures in dollars: Selected countries and years 1960–99	288 289

percent distribution, by type of payer: United States, selected calendar years 1987–2000	298	establishments by size of establishment and occupation: United States, selected years 1990–97	320
120. Employers' costs per employee-hour worked for total compensation, wages and salaries, and health insurance , according to selected characteristics: United States,	200	134. Medicare enrollees and expenditures and percent distribution, according to type of service: United States and other areas, selected years 1970–2000	322
selected years 1991–2001	300	135. Medicare enrollees and program payments among fee-for-service Medicare beneficiaries, according to sex and age: United States and other areas, 1994–99	324
122. Nursing home average monthly charges per resident and percent of residents, according to selected	301	136. Medicare beneficiaries by race and ethnicity, according to selected characteristics: United States, selected years 1992–98	325
facility and resident characteristics: United States, 1977, 1985, 1995, 1997, and 1999	302	137. Medicaid recipients and medical vendor payments, according to basis of eligibility, and race and ethnicity: United States, selected fiscal years 1972–98	327
resident and percent of residents, according to primary source of payments and selected facility characteristics: United States, 1985, 1995, and 1999	303	138. Medicaid recipients and medical vendor payments, according to type of service: United States, selected fiscal years 1972–98	328
124. Mental health expenditures , percent distribution, and per capita expenditures, according to type of mental health organization: United States, selected years 1975–94	304	139. Department of Veterans Affairs health care expenditures and use, and persons treated according to selected characteristics: United States, selected fiscal years	
125. Funding for health research and development, according to source of funds: United States, selected fiscal years 1970–99	305	State Health Expenditures and Health Insura	330 nce
126. Federal spending for human immunodeficiency virus (HIV)-related activities, according to agency and type of activity: United States, selected fiscal years 1985–2001	306	140. Personal health care per capita expenditures, by type of service, geographic division, and State: United States, selected years 1991–98	331
Health Care Coverage and Major Federal Programs		141. Medicare expenditures as a percent of total personal health care expenditures by geographic division and State: United States, 1991–98	337
127. Private health insurance coverage among persons under 65 years of age, according to selected characteristics: United States, selected years 1984–2000	307	142. Medicaid expenditures as a percent of total personal health care expenditures by geographic division and State: United States, 1991–98	338
128. Medicaid coverage among persons under 65 years of age, according to selected characteristics: United States, selected years 1984–2000	310	143. State mental health agency per capita expenditures for mental health services and average annual percent change by geographic division and State: United States,	
129. No health care coverage among persons under 65 years of age, according to selected characteristics: United States, selected years 1984–2000	312	selected fiscal years 1981–97	339
130. Health care coverage for persons 65 years of age and over, according to type of coverage and selected		geographic division and State: United States, 1994 and 1999	340
characteristics: United States, selected years 1989–99 131. Health maintenance organization (HMO) coverage by private insurance, Medicaid, and Medicare according to selected characteristics: United States, 1998, 1999, and	314	145. Medicaid recipients, recipients in managed care, payments per recipient, and recipients per 100 persons below the poverty level by geographic division and State: United States, selected fiscal years 1989–98	341
2000	317	146. Persons enrolled in health maintenance organizations (HMOs) by geographic division and State: United States, selected years 1980–2001	343
	319	147. Persons under 65 years of age without health care coverage by State: United States, selected years 1987–2000	344